

Nucleosynthetic signatures of the first stars

Anna Frebel¹, Wako Aoki², Norbert Christlieb^{2,3}, Hiroyasu Ando², Martin Asplund¹, Paul S. Barklem⁴, Timothy C. Beers⁵, Kjell Eriksson⁴, Cora Fechner³, Masayuki Y. Fujimoto⁶, Satoshi Honda², Toshitaka Kajino², Takeo Minezaki⁷, Ken'ichi Nomoto⁸, John E. Norris¹, Sean G. Ryan⁹, Masahide Takada-Hidai¹⁰, Stelios Tsangarides⁹ & Yuzuru Yoshii⁷

¹Research School of Astronomy & Astrophysics, The Australian National University, Cotter Road, Weston Creek, Australian Capital Territory 2611, Australia

²National Astronomical Observatory of Japan, 2-21-1, Osawa, Mitaka, Tokyo 181-8588, Japan

³Hamburger Sternwarte, Gojenbergsweg 112, 21029 Hamburg, Germany

⁴Department of Physics and Space Sciences, Uppsala Astronomical Observatory, Box 515, SE-751 20 Uppsala, Sweden

⁵Department of Physics and Astronomy, and Joint Institute for Nuclear Astrophysics (JINA), Michigan State University, East Lansing, Michigan 48824, USA

⁶Department of Physics, Hokkaido University, Sapporo, Hokkaido 060-0810, Japan

⁷Institute of Astronomy, School of Science, University of Tokyo, Mitaka, Tokyo 181-0015, Japan

⁸Department of Astronomy, School of Science, University of Tokyo, Tokyo 113-0033, Japan

⁹Department of Physics and Astronomy, Open University, Walton Hall, Milton Keynes MK7 6AA, UK

¹⁰Liberal Arts Education Center, Tokai University, 1117 Kitakaname, Hiratsuka-shi, Kanagawa 259-1292, Japan

The chemically most primitive stars provide constraints on the nature of the first stellar objects that formed in the Universe; elements other than hydrogen, helium and traces of lithium present within these objects were generated by nucleosynthesis in the very first stars. The relative abundances of elements in the surviving primitive stars reflect the masses of the first stars, because the pathways of nucleosynthesis are quite sensitive to stellar masses. Several models^{1–5} have been suggested to explain the origin of the abundance pattern of the giant star HE0107–5240, which hitherto exhibited the highest deficiency of heavy elements known^{1,6}. Here we report the discovery of HE1327–2326, a subgiant or main-sequence star with an iron abundance about a factor of two lower than that of HE0107–5240. Both stars show extreme overabundances of carbon and nitrogen with respect to iron, suggesting a similar origin of the abundance patterns. The unexpectedly low Li and high Sr abundances of HE1327–2326, however, challenge existing theoretical understanding: no model predicts the high Sr abundance or provides a Li depletion mechanism consistent with data available for the most metal-poor stars.

The star HE1327–2326 (coordinates right ascension, RA = 13 h 30 min 06 s and declination, $\delta = -23^{\circ}41'51''$ at Equinox 2000; apparent visual magnitude $V = 13.5$ mag) was found among 1,777 bright ($10 < B < 14$) and apparently metal-poor objects with partly saturated spectra selected from the Hamburg/ESO objective-prism survey (HES)⁷. Its extreme metal-weakness was first appreciated in a medium-resolution ($\delta\lambda \approx 0.5$ nm) follow-up spectrum obtained in April 2003 at the 3.6-m telescope of the European Southern Observatory, Chile. In May and June 2004, observations with the High Dispersion Spectrograph⁸ at the Subaru telescope, Hawaii, were taken. Figure 1 shows a portion of this spectrum and gives more details on the discovery.

An important difference between HE0107–5240 and HE1327–2326 is their evolutionary status. HE0107–5240 is a giant and has evolved off the main-sequence branch and up the red-giant branch.

Consequently, internal mixing has possibly dredged up processed material (such as C, N) from the stellar interior to the surface and altered the abundances of this object. In contrast, HE1327–2326 is relatively unevolved and located on either the main-sequence or the subgiant branch.

We use model atmosphere techniques and the assumption of local thermodynamical equilibrium (LTE) to perform our chemical abundance analysis with non-LTE corrections where available. There are no significant differences between the subgiant and the main-sequence solution. See Table 1 for more details on the analysis procedure and the derived abundances.

HE1327–2326 sets a new record for the star with the lowest iron abundance: $[\text{Fe}/\text{H}]_{\text{non-LTE}} = -5.4 \pm 0.2$ (where $[A/B] = \log_{10}(N_A/N_B) - \log_{10}(N_A/N_B)_{\odot}$ for the number N of atoms of elements A and B and subscript \odot refers to the Sun), derived from four detected weak Fe I lines and corrected for non-LTE. This low value corresponds to $\sim 1/250,000$ of the solar Fe value. To illustrate the nature of such extremely Fe-deficient stars, in Fig. 2 we compare our abundances with those of HE0107–5240 ($[\text{Fe}/\text{H}]_{\text{non-LTE}} = -5.2$;




Figure 1 Comparison of high-resolution spectra of HE1327–2326 with G64–12 and CS 22876–032. The latter is a double-lined spectroscopic binary. All three stars have a similar effective temperature and gravity. In HE1327–2326 we note the absence of the Fe I line at 393 nm together with the appearance of CH lines. Our best fit for the Ca II K ($\lambda = 393.4$ nm) and CH band abundances is plotted in red. Additional CH fits (in red) for our carbon abundance $[\text{C}/\text{Fe}] = 4.1 \pm 0.2$ dex are shown as well. The inset shows our strongest detected Fe I line (0.64 pm equivalent width) at 386 nm in the Subaru data from which we derive $[\text{Fe}/\text{H}]_{\text{non-LTE}} = -5.4 \pm 0.2$. From our medium-resolution spectrum we initially estimated $[\text{Fe}/\text{H}] = -4.0$ for HE1327–2326 using the apparent strength of the Ca II K line index KP of Beers *et al.*²¹ and an approximate colour. The high-resolution data revealed the presence of strong interstellar Ca II K that had not been resolved at the lower resolution. See the prominent feature blueward of the Ca II K line. The interstellar Ca is consistent with a colour excess of $E(B - V) = 0.08$ along the line of sight to HE1327–2326. The presence of CH lines near the Ca II K line in HE0107–5240 caused a similar problem in the initial recognition of its extreme metal-deficiency. These effects should be taken into account in future searches for more stars with similar heavy-element deficiency. Our high-resolution (resolving power $R = \lambda/\Delta\lambda = 50,000$), high signal-to-noise ratio ($S/N = 160$ per pixel at 400 nm) data cover the wavelength range of 304 to 674 nm.

letters to nature

refs 1, 6 and 9), the only other star known to have $[\text{Fe}/\text{H}] < -4.0$. Both objects have very large overabundances of C and N relative to Fe. Although both stars also share a deficiency of Ca and Ti as well as Fe, the Mg and Al abundances relative to Fe in HE1327–2326 are more than one order of magnitude higher than those of HE0107–5240.

Surprisingly, we do not detect the Li I doublet at 670.7 nm in HE1327–2326, although it is found in similarly unevolved metal-poor stars that permit observational determination of the primordial Li abundance and the associated baryon density of the Universe. Our upper limit for Li of $A(\text{Li}) = 1.6$ (where $A(\text{Li}) = \log_{10}(N(\text{Li})/N(\text{H})) + 12.00$) is significantly lower than the primordial Li value of $A(\text{Li}) = 2.62 \pm 0.05$, inferred from the baryon density estimated by the Wilkinson Microwave Anisotropy Probe together with standard Big Bang nucleosynthesis¹⁰. It is even below $A(\text{Li}) \cong 2.0$, found by Ryan *et al.*¹¹ at $[\text{Fe}/\text{H}] = -3.5$. This apparent absence does not result from the evolutionary status of HE1327–2326, because in more evolved metal-poor subgiants such as HD140283, Li has not been depleted. Li weakness occurs rarely in metal-poor stars near the main-sequence turnoff: at effective temperature $T_{\text{eff}} \approx 6,200$ K only 1 in ~ 20 shows this effect¹¹. In HE1327–2326, the weak lines show no evidence for rotational broadening and offer no support for the ultra-Li-depleted rapid rotator model¹², which assumes membership of a binary system, or for rotationally induced mixing in general¹³. Other explanations might include gravitational settling¹⁴ or diffusion, but these have not operated efficiently in other metal-deficient stars of similar temperature and gravity.

Another surprising result is the high Sr abundance (see Fig. 2). Our observed lower limit for the Sr/Ba ratio ($[\text{Sr}/\text{Ba}]_{\text{non-LTE}} > -0.4$) suggests that Sr was not produced in the main ‘slow’-neutron-capture (s-) process occurring in asymptotic giant-branch (AGB) stars, given that the typical ratio in Fe-deficient, C-rich and

s-process-enhanced stars is¹⁵ $[\text{Sr}/\text{Ba}]_{\text{non-LTE}} < -1.0$. The weak s-process occurring in formerly more massive stars (mass $M > 10M_{\odot}$) might yield Sr with little or no Ba, although theoretical calculations suggest that this process is inefficient at low metallicity¹⁶. Alternatively, the main ‘rapid’-neutron-capture (r-) process might have produced the Sr, given that $[\text{Sr}/\text{Ba}]_{\text{non-LTE}} > -0.4$ is consistent with the ratio observed in metal-poor, r-process-enhanced stars, in which¹⁷ $[\text{Sr}/\text{Ba}]_{\text{non-LTE}} \approx -0.3$ (non-LTE corrections were applied as for HE1327–2326). It is therefore possible that the Sr was formed in a type-II supernova, currently believed to be a site of the main r-process that expelled material from which HE1327–2326 formed. Another possibility is the recently proposed process that might have provided light neutron-capture elements in the early Galaxy¹⁶. Although the site and details of this newly suggested process are not yet known, it might provide an explanation for the excesses of light neutron-capture elements, including Sr, seen in very metal-poor stars. To discriminate the contribution of such processes, further constraints on the Ba abundance, as well as further information on other neutron-capture elements, from higher-quality data, are needed.

With its distinctive chemical composition, HE1327–2326 is an ideal tool for ‘stellar archaeology’ and a better understanding of the early enrichment history, and the first mass function of stars in the Universe. Although two stars, HE1327–2326 and HE0107–5240, are now known with $[\text{Fe}/\text{H}] < -5.0$, none has been found in the range $-5 < [\text{Fe}/\text{H}] < -4$. This is suggestive of an underlying, yet to be explored, fundamental physical mechanism in the formation of the first stellar objects² which may differ from that for stars more metal-rich than $[\text{Fe}/\text{H}] = -4.0$.

Table 1 Abundance ratios of HE1327–2326

Element	[X/Fe], A(Li), [Fe/H]	
	Subgiant	Dwarf
Li	<1.6	<1.6
C	4.1 ± 0.2	3.9 ± 0.2
N	4.5 ± 0.2	4.2 ± 0.2
O	<4.0	<3.7
Na (LTE)	2.4 ± 0.2	2.4 ± 0.2
Na (non-LTE)	2.0 ± 0.2	2.0 ± 0.2
Mg (LTE)	1.7 ± 0.2	1.7 ± 0.2
Mg (non-LTE)	1.6 ± 0.2	1.6 ± 0.2
Al (LTE)	1.3 ± 0.2	1.3 ± 0.2
Al (non-LTE)	1.7 ± 0.2	1.7 ± 0.2
Ca I	0.1 ± 0.2	0.1 ± 0.2
Ca II	0.9 ± 0.2	0.8 ± 0.2
Ti	0.6 ± 0.2	0.8 ± 0.2
Fe (LTE)	-5.6 ± 0.2	-5.7 ± 0.2
Fe (non-LTE)	-5.4 ± 0.2	-5.5 ± 0.2
Sr (LTE)	1.0 ± 0.2	1.2 ± 0.2
Sr (non-LTE)	1.1 ± 0.2	1.3 ± 0.2
Ba (LTE)	<1.4	<1.7
Ba (non-LTE)	<1.4	<1.7

Using colour– T_{eff} relations²⁴, we determine $T_{\text{eff}} = 6,180 \pm 80$ K from broad-band UBVR photometry obtained with the MAGNUM telescope²⁵, Hawaii, and JHK magnitudes taken from the 2MASS survey²⁶. To constrain the gravity ($\log g$), we use the proper motion²⁷ ($\mu = 0.073 \pm 0.006$ arcsec yr⁻¹) of HE1327–2326 to set limits on its distance (assuming that the Galactic escape velocity is 500 km s^{-1} and larger than the transverse velocity). It follows that the absolute V magnitude is $M_V > 2.7$ mag which constrains the evolutionary status of the star and therefore $\log g$. Inspection of a 12-billion-year isochrone with $[\text{Fe}/\text{H}] = -3.5$ (the most metal-poor one available²⁸) results in two solutions for the surface gravity $\log g$: 3.7 and 4.5 (in cgs units). Owing to weak observational constraints on $\log g$, it is currently not possible to determine which of these solutions is correct. T_{eff} and $\log g$ are the input for two different plane-parallel LTE model atmospheres that we use to derive the abundances for both gravity solutions; a MARCS model (Gustafsson, B. *et al.*, manuscript in preparation) tailored for the chemical composition of HE1327–2326, and a Kurucz model²⁹. Both sets of abundances agree to within 0.1 dex. Using spectrum synthesis we determine the C and N abundances from CH and NH molecular bands and the upper limit for O from OH molecular bands in the ultraviolet spectral range. Apart from the molecular features, only 13 weak absorption lines are detected in spectral regions not affected by the wings of the Balmer lines and thus suitable for our abundance analysis. Solar abundances were taken from ref. 30. Typical statistical 1σ errors in our abundances are 0.2 dex. Possible systematic errors are judged to be of the same order of magnitude. Non-LTE corrections are included where available²². For the calculation of the non-LTE abundance ratios, the non-LTE iron abundance has been used.




Figure 2 Abundance patterns of HE1327–2326 (subgiant solution, filled circles) and HE0107–5240 (open squares). Typical $1-\sigma$ errors of 0.2 dex are shown in the plot. Upper limits are indicated by an arrow. We adopt the same non-LTE corrections for both stars²², which lead to the modification of the published abundances of HE0107–5240.

Consequently, we adopt $[\text{Fe}/\text{H}]_{\text{non-LTE}} = -5.2$ (ref. 9) as the iron abundance for HE0107–5240. These two most Fe-poor stars both have very large C enhancement relative to Fe by a factor of $\sim 5,000$ (HE0107–5240) and $\sim 10,000$ (HE1327–2326). N/Fe is ~ 30 times the solar value (considering the subgiant solution) in HE1327–2326 whereas it is ~ 200 times the solar value in HE0107–5240. The upper limit for the O abundance of HE1327–2326 is $[\text{O}/\text{Fe}] < 4.0$. Oxygen in HE0107–5240 has recently been determined²³ to be $[\text{O}/\text{Fe}] = 2.3$, which is of the same order as its $[\text{N}/\text{Fe}]$ value. These enormous overabundances in CNO elements suggest that both stars belong to a group of objects sharing a common formation scenario. HE1327–2326 and HE0107–5240 have Ca/Fe and Ti/Fe abundance ratios that are enhanced by factors of less than ten compared to the Sun. The light-element ratios Na/Fe, Mg/Fe and Al/Fe, as well as, surprisingly, Sr/Fe, are all enhanced by factors of ~ 10 to ~ 100 in HE1327–2326. Of these four elements, only Na and Mg are detected in HE0107–5240 with element/Fe ratios close to the solar value. As for several other elements, an upper limit for Ba has been measured in both stars. The Sr/Ba ratio is crucial to identify the origin for the Sr and other heavy elements.

HE1327–2326 permits exploration of the many formation theories of the earliest stars proposed for HE0107–5240. One idea is the pre-enrichment of the gas clouds from which the two stars formed by a first generation ‘faint’ type-II supernova ($M \approx 25M_{\odot}$) experiencing mixing and fallback³, or rotating, massive objects ($M \approx 200\text{--}300M_{\odot}$; ref. 18). In this case, HE1327–2326 would be an early Population II star, formed from gas enriched by one (or possibly a few) of the first type-II supernovae. In particular, the quantitative predictions of abundance patterns by updated models of ‘faint’ type-II supernovae (N. Iwamoto, H. Umeda, N. Tominaga and K. Maeda, personal communication) agree with the chemical abundance patterns of HE1327–2326. The diversity of Mg/Fe ratios found in HE1327–2326 and HE0107–5240 is easily explained by the variety of mixing and fallback. A remaining problem is the Sr, whose production is not yet included in these type-II supernova models.

Another idea is the accretion of heavy elements from the interstellar medium onto first-generation low-mass stars (Population III)^{1,2,19}. In this case, however, the high abundances of lighter elements must be explained by other processes. The unevolved status of HE1327–2326 means that the scenario in which internal mixing processes lead to the dredge-up of processed material can be excluded. A remaining possibility is mass transfer from an AGB star within a binary system^{1,5}. Mass transfer could naturally account for the Li depletion²⁰ found in HE1327–2326. The high Sr abundance is problematic, along with the non-detection of Ba, which cannot be explained by the s-process expected to occur in AGB stars. A crucial test for this scenario is to check the binarity of this object, for which long-period radial velocity monitoring is required. □

Received 22 November 2004; accepted 8 February 2005; doi:10.1038/nature03455.

- Christlieb, N. *et al.* A stellar relic from the early Milky Way. *Nature* **419**, 904–906 (2002).
- Shigeyama, T., Tsujimoto, T. & Yoshii, Y. Excavation of the first stars. *Astrophys. J.* **568**, L57–L60 (2003).
- Umeda, H. & Nomoto, K. First-generation black-hole-forming supernovae and the metal abundance pattern of a very iron-poor star. *Nature* **422**, 871–873 (2003).
- Limongi, M., Chieffi, A. & Bonifacio, P. On the origin of HE 0107–5240, the most iron-deficient star presently known. *Astrophys. J.* **594**, L123–L126 (2003).
- Suda, T., Aikawa, M., Machida, M. N., Fujimoto, M. Y. & Iben, I. Jr. Is HE 0107–5240 a primordial star? The characteristics of extremely metal-poor carbon-rich stars. *Astrophys. J.* **611**, 476–493 (2004).
- Christlieb, N. *et al.* HE 0107–5240, a chemically ancient star. I. A detailed abundance analysis. *Astrophys. J.* **603**, 708–728 (2004).
- Wisotzki, L. *et al.* The Hamburg/ESO survey for bright QSOs. III. A large flux-limited sample of QSOs. *Astron. Astrophys.* **358**, 77–87 (2000).
- Noguchi, K. *et al.* High dispersion spectrograph (HDS) for the Subaru telescope. *Publ. Astron. Soc. Jpn* **54**, 855–864 (2002).
- Beers, T. C. & Christlieb, N. The discovery and analysis of very metal-poor stars in the galaxy. *Annu. Rev. Astron. Astrophys.* (in the press).
- Coc, A., Vangioni-Flam, E., Descouvemont, P., Adahchour, A. & Angulo, C. Updated big bang nucleosynthesis compared with Wilkinson microwave anisotropy probe observations and the abundance of light elements. *Astrophys. J.* **600**, 544–552 (2004).
- Ryan, S. G., Norris, J. E. & Beers, T. C. The Spite lithium plateau: ultrathin but postprimordial. *Astrophys. J.* **523**, 654–677 (1999).
- Ryan, S. G., Gregory, S. G., Kolb, U., Beers, T. C. & Kajino, T. Rapid rotation of ultra-Li-depleted halo stars and their association with blue stragglers. *Astrophys. J.* **571**, 501–511 (2002).
- Pinsonneault, M. H., Walker, T. P., Steigman, G. & Narayanan, V. K. Halo star lithium depletion. *Astrophys. J.* **527**, 180–198 (1999).
- Richard, O., Michaud, G. & Richer, J. Models of metal-poor stars with gravitational settling and radiative accelerations. III. Metallicity dependence. *Astrophys. J.* **580**, 1100–1117 (2002).
- Aoki, W., Norris, J. E., Ryan, S. G., Beers, T. C. & Ando, H. Detection of lead in the carbon-rich, very metal-poor star LP 625–44: A strong constraint on s-process nucleosynthesis at low metallicity. *Astrophys. J.* **536**, L97–L100 (2000).
- Travaglio, C. *et al.* Galactic evolution of Sr, Y, and Zr. A multiplicity of nucleosynthetic processes. *Astrophys. J.* **601**, 864–884 (2004).
- Christlieb, N. *et al.* The Hamburg/ESO R-process enhanced star survey (HERES). I. Project description, and discovery of two stars with strong enhancements of neutron-capture elements. *Astron. Astrophys.* **428**, 1027–1037 (2004).
- Fryer, C. L., Woosley, S. E. & Heger, A. Pair instability supernovae, gravity waves, and gamma-ray transients. *Astrophys. J.* **550**, 372–382 (2001).
- Yoshii, Y. Metal enrichment in the atmospheres of extremely metal-deficient dwarf stars by accretion of interstellar matter. *Astron. Astrophys.* **97**, 280–290 (1981).
- Norris, J. E., Ryan, S. G., Beers, T. C. & Deliyannis, C. P. Extremely metal-poor stars. III. The Li-depleted main-sequence turnoff dwarfs. *Astrophys. J.* **485**, 370–379 (1997).
- Beers, T. C., Rossi, S., Norris, J. E., Ryan, S. G. & Sheffer, T. Estimation of stellar metal abundance. II. A recalibration of the Ca II K technique, and the autocorrelation function method. *Astron. J.* **117**, 981–1009 (1999).
- Asplund, M. New light on stellar abundances analyses: departures from LTE and homogeneity. *Annu. Rev. Astron. Astrophys.* (in the press).
- Bessell, M. S., Christlieb, N. & Gustafsson, B. On the oxygen abundance of HE 0107–5240. *Astrophys. J.* **612**, L61–L63 (2004).
- Alonso, A., Arribas, S. & Martinez-Roger, C. The empirical scale of temperatures of the low main sequence (F0V–K5V). *Astron. Astrophys.* **313**, 873–890 (1996).
- Yoshii, Y. in *New Trends in Theoretical and Observational Cosmology* (eds Sato, K. & Shiromizu, T.) 235–244 (Universal Academy, Tokyo, 2002).
- Cutri, R. M., *et al.* 2MASS All-Sky Catalog of Point Sources (California Institute of Technology, Pasadena, 2003); (<http://irsa.ipac.caltech.edu/applications/Gator>).
- Girard, T. M. *et al.* The southern proper motion program. III. A near-complete catalog to $V = 17.5$. *Astron. J.* **127**, 3060–3071 (2004).
- Kim, Y., Demarque, P., Yi, S. K. & Alexander, D. R. The Y2 isochrones for alpha-element enhanced mixtures. *Astrophys. J. Suppl.* **143**, 499–511 (2002).
- Kurucz, R. L. *ATLAS9 Stellar Atmosphere Programs and 2 km/s Grid* CD-ROM 13 (Smithsonian Astrophysical Observatory, Cambridge, 1993); (<http://kurucz.harvard.edu/cdroms.html>).
- Asplund, M., Grevesse, N. & Sauval, A. J. in *Cosmic Abundances As Records Of Stellar Evolution And Nucleosynthesis* (eds Bash, F. N. & Barnes, T. G.) *ASP Conf. Ser.* (in the press); preprint at (<http://www.arxiv.org/astro-ph/0410214>) (2004).

Acknowledgements We thank A. Steinhauer and C. Thom for obtaining additional observations, N. Iwamoto, K. Maeda, T. Suda, N. Tominaga and H. Umeda for valuable discussions and L. Wisotzki and D. Reimers for leading the HES. This work was supported by the Astronomical Society of Australia (A.F.), Australian Research Council (M.A., A.F., J.E.N.), Ministry of Education, Culture, Sports, Science and Technology in Japan and JSPS (all Japanese co-authors), Deutsche Forschungsgemeinschaft (N.C.), Swedish Research Council (P.S.B., K.E.), US National Science Foundation (T.C.B.) and JINA (T.C.B., N.C., A.F., J.E.N.). This work is based on data collected at the Subaru Telescope, which is operated by the National Astronomical Observatory of Japan.

Competing interests statement The authors declare that they have no competing financial interests.

Correspondence and requests for materials should be addressed to A.F. (anna@mso.anu.edu.au).